

# Notes: Bernoulli, Binomial, and Geometric Distributions

CS 3130/ECE 3530: Probability and Statistics for Engineers

September 19, 2017

## **Bernoulli distribution:**

Defined by the following pmf:

$$p_X(1) = p, \quad \text{and} \quad p_X(0) = 1 - p$$

Don't let the  $p$  confuse you, it is a single number between 0 and 1, not a probability function. If  $X$  is a random variable with this pmf, we say “ $X$  is a Bernoulli random variable with parameter  $p$ ”, or we use the notation  $X \sim Ber(p)$ . You can think of a Bernoulli trial as flipping a coin where the chance of heads is  $p$  and the chance of tails is  $1-p$ . Often we call 0 a “failure” and 1 a “success”, so  $p$  is the probability of success.

## **Binomial distribution:**

The binomial distribution describes the probabilities for repeated Bernoulli trials – such as flipping a coin ten times in a row. Each trial is assumed to be independent of the others (for example, flipping a coin once does not affect any of the outcomes for future flips). First, we need some definitions.

Remember the definition for factorial:

$$n! = n \times (n - 1) \times \cdots \times 2 \times 1$$

This is the number of ways to put  $n$  objects into a specific order.

And the definition for “ $n$  choose  $k$ ”:

$$\binom{n}{k} = \frac{n!}{(n - k)! k!}$$

This is the number of ways to select  $k$  objects out of a possible  $n$ , where the order does *not* matter.

The **binomial distribution** with parameters  $n$  and  $p$  is given by the pmf:

$$p_X(k) = \binom{n}{k} p^k (1 - p)^{n-k}.$$

This is denoted  $X \sim Bin(n, p)$ . This distribution is for repeated Bernoulli trials, and it gives the probability that you get  $k$  successes out of  $n$  trials.

## **Geometric distribution:**

The **geometric distribution** is also for repeated Bernoulli trials, and it gives the probability that the first  $k - 1$  trials are failures, while the  $k$ th trial is the first success. Its pmf is

$$p_X(k) = (1 - p)^{k-1} p.$$

This is denoted  $X \sim Geo(p)$ .

In-class Problem: Remember the Monty Hall problem – if we switch doors, we have a  $2/3$  chance of winning and  $1/3$  chance to lose. If we play the game 4 times, what is the probability that we win exactly once? How about exactly 0, 2, 3, or 4 times? What is the chance that we lose the first three times and finally win on the 4th try?

**Key to variable names**

It's important to keep straight what all the variables mean in the above equations. Here is a summary:

$n$  : Number of trials

$k$  : Number of successes in Binomial, OR first success that occurs in Geometric

$p$  : Probability of success